

Data-driven Estimation of SM Backgrounds for SUSY Searches at the LHC

Takayuki Yamazaki,
on behalf of ATLAS and CMS collaborations
Graduate School of Science, University of Tokyo
7-3-1 Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

Searches for supersymmetry (SUSY) are very important tasks at the Large Hadron Collider (LHC). If SUSY exists at the TeV scale, clear excess above the Standard Model (SM) background will be observed. SM background should be estimated from real data self. In this paper, we describe the strategy for the early SUSY searches at the LHC and focus on the data-driven estimation of the SM background in the early stage of collision.

1 Introduction

The Standard Model (SM) can explain almost all experimental results so far but is not considered to be a complete theory. Supersymmetry (SUSY), which is symmetry between fermions and bosons, is the most attractive theoretical model for new physics beyond the SM. If SUSY particles exist at the TeV scale, the hierarchy problem can be resolved and three forces, electromagnetic, weak and strong interaction are unified at the GUT scale ($\sim 10^{16}\text{GeV}$). Moreover, the lightest SUSY particle (LSP) is a good candidate for the cold dark matter if R-parity is conserved.

At the Large Hadron Collider (LHC), which will start to operate in 2008. Direct searches of the physics at the TeV scale can be performed. The strategy for early SUSY searches is described in this paper, and I focus on the estimation of SM background using the data corresponding an integrated luminosity of 1fb^{-1} .

2 SUSY Signatures

Sparticles such as gluinos ($\tilde{g}$) and squarks ($\tilde{q}$) will be produced via strong interaction at the LHC. They decay into the lightest state of the SUSY particle and SM particles. The lightest SUSY particle is stable if R-parity is conserved. Final event topology is high p_T multijets and large missing E_T , sometimes including leptons. This is model-independent signature of the SUSY events, and our SUSY search is based on this experimental signature.

Search modes are classified by the number of isolated leptons. The promising search modes are listed in Table 1. They cover realistic SUSY models such as minimal supergravity (mSUGRA), gauge mediated SUSY breaking (GMSB), and anomaly mediated SUSY breaking (AMSB).

The selection criteria of SUSY searches are

- 4 jets with $p_T > 50 \text{ GeV}/c$ and at least 1 jet with $p_T > 100 \text{ GeV}/c$
- $S_T > 0.2$

Table 1: Search modes, corresponding SUSY models and SM background processes.

number of leptons	SUSY model	SM background
no lepton	mSUGRA, AMSB	QCD, $t\bar{t}$, W , Z
one lepton (e, μ)	mSUGRA, AMSB	$t\bar{t}$, W
dilepton (e, μ)	mSUGRA, AMSB, GMSB	$t\bar{t}$

- $E_T^{miss} > 100$ GeV and $E_T^{miss} > 0.2 \times M_{eff}$,

where S_T is the transverse sphericity and M_{eff} is the effective mass. M_{eff} is calculated from the leading four jets, E_T^{miss} , and the leptons.

$$M_{eff} = \sum_{i=1}^4 p_T^{jeti} + E_T^{miss} (+ \sum_j p_T^{leptonj}). \quad (1)$$

After the selection, the SUSY signal with large E_T^{miss} can be observed as an excess from the SM background. Figure 1 shows E_T^{miss} distribution for no lepton search mode. The signal shows the promising SUSY point, where $m_0 = 100$ GeV, $m_{1/2} = 300$ GeV, $A = 300$ GeV, $\tan\beta = 6$ and $\mu > 0$. Clear excess over the SM background can be observed with an integrated luminosity of 1fb^{-1} .




Figure 1: The E_T^{miss} distributions for no lepton (left) and one lepton (right) mode. The histograms are normalized to the integrated luminosity of 1fb^{-1} .

3 Data-drive Estimation of the Standard Model Backgrounds

It is essential to understand and estimate the SM background for concrete evidence of SUSY. We take data-driven approach to estimate the SM background because the uncertainties of Monte Carlo predictions are expected to be large in the early stage of the experiment.

3.1 M_T Method

Top pair ($t\bar{t}$) and $W^\pm + \text{jets}$ processes are the dominant backgrounds for one lepton SUSY search mode (Fig. 1). For one lepton mode, $M_T > 100$ GeV is required to enhance the SUSY signal (Fig. 2). M_T is the transverse mass between the lepton and missing energy. The control sample with $M_T < 100$ GeV is used for the estimation. Since SUSY contribution in the control sample is much smaller than in the signal region and the shapes of distributions such as E_T^{miss} are

independent of M_T , the SM background in the signal region can be estimated from the control sample as shown in Figure 2.




Figure 2: The M_T distribution (left) and the estimation of E_T^{miss} distribution of the SM background (right) for one lepton SUSY search mode. The histograms are normalized to the integrated luminosity of 1fb^{-1} .

3.2 QCD Background

The QCD processes contribute to the backgrounds of no lepton mode (Fig. 1). In order to reduce the QCD background, we add one more cut which require the difference in azimuthal angle ϕ between E_T^{miss} and the three most energetic jets to be larger than 0.2. The distributions of the minimal value of the $\Delta\phi$ (min $\Delta\phi$) are shown in Figure 3 and the min $\Delta\phi$ distribution of the QCD processes has small $\Delta\phi$ because E_T^{miss} in the QCD processes is due to neutrinos emitted from semileptonic decays of b/c quarks or mis-measurement of jet energies.

The QCD background can be estimated from multi-jet events without E_T^{miss} . The missing E_T function, which is the momentum fraction of the neutrino in b/c decay, is applied to the 1st and 2nd leading jet of the control sample and then the heavy-flavor QCD events with large E_T^{miss} can be reproduced. Figure 3(b) shows E_T^{miss} distributions of true QCD processes and estimation with this method.




Figure 3: The minimal $\delta\phi$ distribution (left) and the estimation of E_T^{miss} distribution of the QCD events (right) in no lepton SUSY search mode. The histograms are normalized to the integrated luminosity of 1fb^{-1} .

4 Conclusion

The LHC will start to operate in 2008 and the data-driven method can be used for the background estimation of the SUSY searches in the early stage of the collision. The essence of the

data driven approach are summerized in this paper. Our searches are based on the SUSY event signitures and less model independent but we can discover SUSY whose squarks and gluino masses up to 1 TeV with an integrated luminosity of 1fb^{-1} .

Acknowledgments

The author would like to thank all members of ATLAS and CMS collaborations.

References

1. The ATLAS collaboration, *ATLAS Detector and Physics Performance Technical Design Report*, CERN/LHCC/99-14 and CERN/LHCC/99-15.
2. The CMS collaboration, *CMS Physics Technical Design Report*, CERN/LHCC/2006-001 and CERN/LHCC/2006-021.